Exposure Dose Distribution of the Workers at Fukushima Daiichi Nuclear Power Plant

(Updated on 30 September 2019)

1 Radiation Exposure Dose Distributions

(1) The distribution of external exposure dose of the workers during the last 3 months (Numbers of workers who entered each area every month)

(rumbers of workers who entered each area every month)									
Effective dose (E)	June 2019			July 2019			August 2019		
mSv	TEPCO	Contractors	Total	TEPCO	Contractors	Total	TEPCO	Contractors	Total
100 <e< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></e<>	0	0	0	0	0	0	0	0	0
75 <e≤100< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></e≤100<>	0	0	0	0	0	0	0	0	0
50 <e≤75< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></e≤75<>	0	0	0	0	0	0	0	0	0
20 <e≤50< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></e≤50<>	0	0	0	0	0	0	0	0	0
10 <e≤20< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></e≤20<>	0	0	0	0	0	0	0	0	0
5 <e≤10< td=""><td>1</td><td>12</td><td>13</td><td>0</td><td>10</td><td>10</td><td>0</td><td>4</td><td>4</td></e≤10<>	1	12	13	0	10	10	0	4	4
1 <e≦5< td=""><td>17</td><td>538</td><td>555</td><td>19</td><td>547</td><td>566</td><td>12</td><td>467</td><td>479</td></e≦5<>	17	538	555	19	547	566	12	467	479
E≤1	1,016	4,993	6,009	964	5,048	6,012	1,005	5,037	6,042
Total	1,034	5,543	6,577	983	5,605	6,588	1,017	5,508	6,525
Maximum (mSv)	5.20	7.11	7.11	3.60	9.70	9.70	2.75	7.60	7.60
Average (mSv)	0.12	0.35	0.31	0.13	0.35	0.32	0.09	0.28	0.25

(*) Exposure doses and the number of workers are subject to change due to the replacement of accumulated doses measured using PAD with monthly doses measured using an integrating dosimeter and the reflection of values for workers wearing only an integrating dosimeter (e.g., workers working only within a seismically isolated building).

(2) Combined Cumulative Effective Dose from April 2016 (Internal and External)

Effective dose (E)	April	2016 - July	2019	April 2016 - August 2019 Differen					
mSv	TEPCO	Contractors	Total	TEPCO	Contractors	Total	TEPCO	Contractors	Total
100 <e< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></e<>	0	0	0	0	0	0	0	0	0
75 <e≤100< td=""><td>0</td><td>7</td><td>7</td><td>0</td><td>9</td><td>9</td><td>0</td><td>2</td><td>2</td></e≤100<>	0	7	7	0	9	9	0	2	2
50 <e≤75< td=""><td>0</td><td>112</td><td>112</td><td>0</td><td>126</td><td>126</td><td>0</td><td>14</td><td>14</td></e≤75<>	0	112	112	0	126	126	0	14	14
20 <e≤50< td=""><td>41</td><td>1,509</td><td>1,550</td><td>45</td><td>1,518</td><td>1,563</td><td>4</td><td>9</td><td>13</td></e≤50<>	41	1,509	1,550	45	1,518	1,563	4	9	13
10 <e≤20< td=""><td>138</td><td>2,106</td><td>2,244</td><td>135</td><td>2,127</td><td>2,262</td><td>-3</td><td>21</td><td>18</td></e≤20<>	138	2,106	2,244	135	2,127	2,262	-3	21	18
5 <e≤10< td=""><td>171</td><td>2,224</td><td>2,395</td><td>175</td><td>2,228</td><td>2,403</td><td>4</td><td>4</td><td>8</td></e≤10<>	171	2,224	2,395	175	2,228	2,403	4	4	8
1 <e≦5< td=""><td>560</td><td>4,485</td><td>5,045</td><td>560</td><td>4,507</td><td>5,067</td><td>0</td><td>22</td><td>22</td></e≦5<>	560	4,485	5,045	560	4,507	5,067	0	22	22
E≤1	1,307	8,801	10,108	1,324	8,850	10,174	17	49	66
Total	2,217	19,244	21,461	2,239	19,365	21,604	22	121	143
Maximum (mSv)	38.14	79.90	79.90	38.09	79.90	79.90	-	-	-
Average (mSv)	2.66	5.88	5.54	2.68	5.92	5.58	-	-	-

(*) Exposure doses and the number of workers are subject to change due to the replacement of accumulated doses measured using PAD with monthly doses measured using an integrating dosimeter and the reflection of values for workers wearing only an integrating dosimeter (e.g., workers working only within a seismically isolated building).

Effective dose (E)	April	2019 - July	2019	April 2	019 - Augus	t 2019			
mSv	TEPCO	Contractors	Total	TEPCO	Contractors	Total	TEPCO	Contractors	Total
100 <e< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></e<>	0	0	0	0	0	0	0	0	0
75 <e≤100< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></e≤100<>	0	0	0	0	0	0	0	0	0
50 <e≤75< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></e≤75<>	0	0	0	0	0	0	0	0	0
20 <e≤50< td=""><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></e≤50<>	0	0	0	0	0	0	0	0	0
10 <e≤20< td=""><td>0</td><td>52</td><td>52</td><td>1</td><td>80</td><td>81</td><td>1</td><td>28</td><td>29</td></e≤20<>	0	52	52	1	80	81	1	28	29
5 <e≦10< td=""><td>10</td><td>332</td><td>342</td><td>13</td><td>445</td><td>458</td><td>3</td><td>113</td><td>116</td></e≦10<>	10	332	342	13	445	458	3	113	116
1 <e≤5< td=""><td>144</td><td>1,425</td><td>1,569</td><td>173</td><td>1,599</td><td>1,772</td><td>29</td><td>174</td><td>203</td></e≤5<>	144	1,425	1,569	173	1,599	1,772	29	174	203
E≤1	1,093	5,062	6,155	1,102	5,044	6,146	9	-18	-9
Total	1,247	6,871	8,118	1,289	7,168	8,457	42	297	339
Maximum (mSv)	9.30	18.30	18.30	10.85	18.30	18.30	-	-	-
Average (mSv)	0.41	1.08	0.98	0.47	1.25	1.13	-	-	-

(3) Combined Cumulative Effective Dose from April 2019 (Internal and External)

(*) Exposure doses and the number of workers are subject to change due to the replacement of accumulated doses measured using PAD with monthly doses measured using an integrating dosimeter and the reflection of values for workers wearing only an integrating dosimeter (e.g., workers working only within a seismically isolated building).

(4) Distribution of sum of external exposure dose and internal exposure dose of workers engaged in specified high-dose work*

(Specified high-dose work has not been performed since October 2015.)

Effective dose (E) mSv	March 2011- September 2015					
100 <e< td=""><td>1</td></e<>	1					
75 <e≤100< td=""><td>191</td></e≤100<>	191					
50 <e≤75< td=""><td>233</td></e≤75<>	233					
20 <e≤50< td=""><td colspan="6">267</td></e≤50<>	267					
10 <e≤20< td=""><td colspan="6">186</td></e≤20<>	186					
5 <e≤10< td=""><td colspan="6">129</td></e≤10<>	129					
1 <e≤5< td=""><td>145</td></e≤5<>	145					
E≤1	51					
Total	1,203					
Maximum (mSv)	102.69					
Average (mSv)	36.49					

(*) Workers engaged in work to which dose limit (100 mSv) during emergency work is applied in line with Article 7 of the Ordinance on Prevention of Ionizing Radiation Hazards.

Specifically, these workers are those who are engaged in work to maintain the functions of a nuclear reactor facility or spent fuel storage pool, or in work to maintain functions to suppress or prevent the possible release of a large amount of radioactive materials due to a failure of or damage to the nuclear reactor facility at a location around the nuclear reactor facility, steam turbine, or accessory facility where hourly dose may exceed 0.1 mSv. It should be noted that only TEPCO employees have so far been engaged in specified high-dose work.

- (*) Workers engaged in specified high-dose work in each month is the number of workers registered as workers engaged in specified high-dose work in that month.
- However, the total of March 2011 to September 2015 includes workers released from specified high-dose work.
- (*) Exposure doses and the number of workers are subject to change due to the replacement of accumulated doses measured using PAD with monthly doses measured using an integrating dosimeter and the reflection of values for workers wearing only an integrating dosimeter (e.g., workers working only within a seismically isolated building).
- (*) The results of re-evaluating committed doses in March 2011 reveal that maximum cumulative effective doses for the period between March 2011 and September 2015 exceeded 100.